

# Tiling of regular polygon with similar right triangles, II

Vigdorichik Leonid\*

A tiling is a decomposition of a polygon into finitely many non-overlapping triangles. (Note that a tiling is not a triangulation, i.e. a vertex of a triangle can lie on a side of a triangle.)

**Theorem 1.** *If a regular  $n$ -gon,  $n > 8$ ,  $n \neq 12, 14, 20, 32, 44$ , can be tiled with similar right triangles, then no angle of this triangle equals to  $\frac{n-2}{3n}\pi$ .*

Theorem 1 follows from [L20, Theorem 1] for all  $n \geq 25$ ,  $n \neq 32, 42$ .

**Corollary 2.** *A regular  $n$ -gon,  $n > 8$ ,  $n \neq 12, 14, 20, 28, 32, 44$ , can be tiled by similar right triangles then one of the angles of these triangles is in  $\frac{\pi}{n}$  or  $\frac{2\pi}{n}$ .*

Theorem 1 and [V. Theorem 1] imply Corollary 2.

We denote  $a_n = 2\frac{n-2}{3n}$  for integer  $n$ . Then  $1 - a_n = \frac{n+4}{3n}$ .

For all  $n$ :  $a_n < 2/3$  and  $1 - a_n > 1/3$ .

**Lemma 3.** *If  $3a_n = p(1 - a_n) + qa_n + r$ ,  $n > 8$ ,  $n \neq 14, 20, 32$ , and  $p, q, r$  are non-negative integers, then  $q = 3$ .*

*Proof.* Since  $\frac{r}{3} \leq a_n = 2\frac{n-2}{3n} < \frac{2}{3}$ , it follows that  $r < 2$ . Since  $qa_n \leq 3a_n$ , it follows that  $q \leq 3$ . We also have  $p = \frac{a_n(3-q)-r}{1-a_n}$ ,  $1 - a_n > \frac{1}{3}$ ,  $a_n < \frac{2}{3}$  therefore  $p < \frac{\frac{2}{3}(3-q)-r}{\frac{1}{3}} = 2(3-q) - 3r$ . We consider all possible values  $r, q$  and show that either  $q = 3$  or the values of  $n$  that correspond to the integer values of  $p$  that do not suit to limitations of the lemma:

1. Let  $r = 0$ .
  - 1.1 Let  $q = 0$ . This is only possible if  $0 \leq p < 6$ . Then  $n \in \{2, 5, 8, 14, 32\}$ .
  - 1.2 Let  $q = 1$ . This is only possible if  $0 \leq p < 4$ . Then  $n \in \{2, 4, 8, 20\}$ .
  - 1.3 Let  $q = 2$ . This is only possible if  $0 \leq p < 2$ . Then  $n \in \{2, 8\}$ .
  - 1.4 Let  $q = 3$ . Then we get the statement of the lemma.
2. Let  $r = 1$ . Since  $\frac{1}{3} < a_n < \frac{2}{3}$ , it follows that  $q \leq \frac{3a_n-1}{a_n} < 2$ .
  - 2.1 Let  $q = 0$ . This is only possible if  $0 \leq p < 3$ . Then  $n \in \{2, 4, 8, 20\}$ .
  - 2.2 Let  $q = 1$ . This is only possible if  $0 \leq p < 1$ . Then  $n = 8$ . □

**Lemma 4.** *If  $b = 1, 2, 3, 4$  and  $b = p(1 - a_n) + qa_n + r$  where  $n > 8$ ,  $n \neq 12, 14, 20, 32, 44$  and  $p, q, r$  are non-negative integers, then  $q \geq p$ .*

\*This work was prepared in frame of math circle "Math and Olympiades".

*Proof.* If  $r = 1$  then  $p, q = 1$ . If  $1 = p(1 - a_n) + qa_n + r$ , then if  $q = 0$  we have  $p = \frac{1}{1-a_n}$ . But  $\frac{1}{3} < 1 - a_n < \frac{1}{2}$ . Hence  $p$  is not non-negative integer. Contradiction. Hence  $p, q = 1$ .

If  $r > 0$  we have the previous case. If  $2 = p(1 - a_n) + qa_n + r$ , then if  $q = 1$  we have the previous case. Hence  $p = \frac{2}{1-a_n} = \frac{6n}{n+4}$ . Then  $0 < \frac{6n}{n+4} < 7$  is non-negative integer. Hence  $n = 20$ . Hence  $p, q = 2$ .

If  $r > 0$  we have one of the previous cases. If  $3 = p(1 - a_n) + qa_n + r$ , then if  $q = 1$  we have the previous case. Hence  $p = \frac{3}{1-a_n} = \frac{9n}{n+4}$ . Then  $0 < \frac{9n}{n+4} < 10$  is non-negative integer. Hence  $n \in \{14, 32\}$ . Hence  $p, q = 3$ .

If  $r > 0$  we have one of the previous cases. If  $4 = p(1 - a_n) + qa_n + r$ , then if  $q = 1$  we have the previous case. Hence  $p = \frac{4}{1-a_n} = \frac{12n}{n+4}$ . Then  $0 < \frac{12n}{n+4} < 13$  is non-negative integer. Hence  $n \in \{12, 20, 44\}$ . Hence  $p, q = 4$ .  $\square$

*Proof of Theorem 1.* Assume that a regular  $n$ -gon,  $n > 8$ ,  $n \neq 12, 14, 20, 32, 44$ , is tiled with similar right triangles of angles  $\frac{\pi}{6} < \alpha < \frac{\pi}{4}$ ,  $\frac{\pi}{2} - \alpha$  and  $\frac{\pi}{2}$ , and  $\alpha = \frac{n+4}{6n}\pi$ .

Take a vertex of the  $n$ -gon. By  $p, q, r$  we denote the number of smaller acute angles  $\alpha$ , bigger acute angles  $\frac{\pi}{2} - \alpha$  and angles  $\frac{\pi}{2}$  respectively at this vertex. Then  $3a_n = p(1 - a_n) + qa_n + r$ , where  $\frac{1}{2} < 3a_n < \frac{2}{3}$ . So  $a_n = 2\frac{\frac{\pi}{2} - \alpha}{\pi}$ ,  $1 - a_n = 2\frac{\alpha}{\pi}$ ,  $r = 2\frac{\frac{\pi}{2} - \alpha}{\pi}$ . Then by Lemma 3 we have  $q = 3, p = 0$ .

For the triangles that have same vertices inside of regular  $n$ -gon or that have same vertices on the side of  $n$ -gon or on the side of a triangle we denote  $p, q, r$  as in previous case. Then by Lemma 4  $p \leq q$ .

Hence the number of bigger acute angles is greater than the number of smaller acute angles. Hence our tiling is not realizable.  $\square$

[V] I.Vasenov, Tiling of regular polygon with similar right triangles,  
<https://arxiv.org/abs/2010.05052>

[L20] M.Laczkovich, Irregular tilings of regular polygons with similar triangles,  
<https://arxiv.org/abs/2002.12013>