

Tiling of regular polygon with similar right triangles I

Vasenov Ivan*

A *tiling* is a decomposition of a polygon into finitely many non-overlapping triangles. (Note that a tiling is not a triangulation, i.e. a vertex of a triangle can lie on a side of a triangle.)

Conjecture 1. *If a regular n -gon, $n \geq 5$, $n \neq 28$, can be tiled with similar right triangles, then one of the angles of these triangles is in $\{\frac{\pi}{n}, \frac{2\pi}{n}, \frac{\pi}{6} + \frac{2\pi}{3n}\}$.*

From [L20, Theorem 1] it follows that if a regular n -gon, $n \geq 25$, $n \neq 30, 42$ is tiled with similar right triangles with angles $\alpha, \frac{\pi}{2} - \alpha, \frac{\pi}{2}$, $\alpha < \frac{\pi}{4}$ then $\alpha = \frac{\pi}{n}$ (see Remark 8 (d)).

Let us present a simple deduction of the corollary from [L20, Theorem 1]. Suppose that at the vertex of the regular n -gon there are p angles α , q angles $\frac{\pi}{2} - \alpha$ and r angles $\frac{\pi}{2}$. From Lemma 6 it follows that α is a rational multiple of π , which satisfies the conditions of [L20, Theorem 1].

Conjecture 2. *If a regular 8-gon can be tiled with similar right triangles, then one of the angles of these triangles is in $\{\frac{\pi}{8}, \frac{\pi}{4}\}$.*

Conjecture 1 and [V. Theorem 1] imply the following corollary.

Conjecture 3. *A regular n -gon, $9 \leq n$, $n \neq 12, 14, 20, 32, 44$, can be tiled by similar right triangles then one of the angles of these triangles is in $\frac{\pi}{n}$ or $\frac{2\pi}{n}$.*

Clearly, for each n there is the ‘trivial’ tiling of a regular n -gon obtained by joining the center of the n -gon to each vertex of the n -gon. We obtain n congruent isosceles triangles with angles $\frac{2\pi}{n}$ at their vertices opposite to their bases. The ‘trivial’ tiling with isosceles triangles gives a ‘trivial’ tiling with congruent right triangles having angles $\frac{\pi}{n}$. We do not know if the remaining value $\frac{2\pi}{n}$ from Theorem 1 are ‘realizable’ by some tilings for $7 \leq n \leq 24$ or $n = 30, 42$.

For $5 \leq n \leq 24$ or $n \in 30, 42$ conjecture 1 cannot be obtained from [L20, Theorem 1] and [L12, Theorem 2.1] (see Remark 8 (e)).

Our proof of conjecture 1 is based on showing that for angles of the triangles other than mentioned in statement *the number of smaller acute angles* is greater

*This work was prepared in frame of math circle ”Math and Olympiads”. High School 67, Kutuzovsky prospect, 10, Moscow, Russia.

than the *number of larger acute angles*. This idea reduces the amount of cases to search through.

We illustrate the idea by a proof of the conjecture 2. If a reader is only interested in the proof of conjecture 1, he could skip conjecture 4 and the proof of conjecture 2.

By the side of a triangle we mean side of a triangle without its vertex.

Conjecture 4. *If $\frac{3}{2} = pa + q(1 - a) + r$, where $0 < a < \frac{1}{2}$, $a \neq \frac{1}{4}$ and p, q, r are non-negative integers, then $p > q$ and $a = \frac{3}{2s}$ for some positive integer s .*

Sketch of proof of conjecture 4. Assume to the contrary that $q - p \geq 0$. We have

$$\frac{3}{2} = pa + q(1 - a) + r = (q - p)(1 - a) + p + r.$$

If $q - p = 0$ then $\frac{3}{2} = p + r$, but $p + r$ is an integer. Then $q - p \geq 1$.

Since $\frac{1}{2} < 1 - a = \frac{\frac{3}{2} - p - r}{q - p} \leq \frac{3}{2} - p - r$, we have $r = p = 0$.

Thus $\frac{3}{2} = (1 - a)q$. $q \neq 0$ since $\frac{3}{2} \neq 0$. Since $0 < a < \frac{1}{2}$, it follows that $\frac{q}{2} < \frac{3}{2} < q$. Then $q = 2$. Hence $a = \frac{1}{4}$. A contradiction.

Thus $p > q$. We have

$$\frac{3}{2} = pa + q(1 - a) + r = (p - q)a + q + r.$$

Since $0 < a$ and q, r are non-negative integers, it follows that $0 \leq q + r < 2$. Then $q + r \in \{0, 1\}$. Hence $a = \frac{3 - 2(q + r)}{2(p - q)} \in \{\frac{3}{2(p - q)}, \frac{1}{2(p - q)}\}$.

Thus $a = \frac{3}{2s}$, where s is a positive integer. \square

Conjecture 5. *If $4 = pa + q(1 - a) + r$, where $0 < a < \frac{1}{2}$, $a \notin \{\frac{1}{4}, \frac{1}{5}, \frac{2}{5}, \frac{3}{7}, \frac{1}{3}\}$, p, q, r are non-negative integers, then $p \geq q$.*

Proof. Assume to the contrary that $q - p > 0$. Then $q - p \geq 1$. We have

$$4 = pa + q(1 - a) + r = (q - p)(1 - a) + r + p.$$

If $r + p > 2$, then $1 - a = \frac{4 - p - r}{q - p}$ is either greater than or equal to 1 or is less than or equal to $\frac{1}{2}$.

If $r + p = 2$, then $1 - a = \frac{2}{q - p}$. Since $0 < a < \frac{1}{2}$ it follows that $a = \frac{1}{3}$.

If $r + p = 1$, then $1 - a = \frac{3}{q - p}$. Hence $a \in \{\frac{1}{4}, \frac{2}{5}\}$.

If $r + p = 0$, then $1 - a = \frac{4}{q - p}$. Hence $a \in \{\frac{1}{5}, \frac{1}{3}, \frac{3}{7}\}$.

A contradiction. Thus, $p \geq q$. \square

Sketch of proof of conjecture 2 independent of conjecture 1. Suppose there is a tiling of a regular 8-gon with right triangles of angles $\alpha < \frac{\pi}{4}$, $\frac{\pi}{2} - \alpha$ and $\frac{\pi}{2}$, and $\alpha \neq \frac{\pi}{8}$.

Take a vertex of the 8-gon. By p, q, r we denote the number of smaller acute angles α , bigger acute angles $\frac{\pi}{2} - \alpha$ and angles $\frac{\pi}{2}$ at this vertex, respectively. Then $\frac{3\pi}{4} = p\alpha + q(\frac{\pi}{2} - \alpha) + r\frac{\pi}{2}$. Divide this equality by $\frac{\pi}{2}$ and get $\frac{3}{2} = pa + q(1 - a) + r$, where $0 < a < \frac{1}{2}$.

Then by conjecture 4

- at each vertex of an 8-gon the number of smaller acute angles α is greater than the number of bigger acute angles $\frac{\pi}{2} - \alpha$.

- $\alpha = \frac{3\pi}{4s}$ for some positive integer s .

Hence $\alpha \notin \{\frac{\pi}{10}, \frac{\pi}{5}, \frac{3\pi}{14}, \frac{\pi}{6}\}$.

For the triangles that have same vertices inside of the 8-gon and not on the side of a triangle we use conjecture 5. Then at any point inside of the 8-gon and not on the side of a triangle the number of smaller acute angles is greater than or equal to the number of bigger acute angles.

For the triangles that have same vertices on the side of 8-gon or on the side of a triangle, we use conjecture 5, substituting $r - 2$ for r . Then at any point on the side of 8-gon or on the side of a triangle the number of smaller acute angles is greater or equal to the number of bigger acute angles.

Hence the number of smaller acute angles is greater than the number of bigger acute angles. A contradiction. \square

Conjecture 6. For any integer $n \geq 5$ if $2 - \frac{4}{n} = pa + q(1 - a) + r$, where $0 < a \leq \frac{1}{2}$, $a \notin \{\frac{2}{n}, \frac{4}{n}, \frac{1}{3} + \frac{4}{3n}\}$, p, q, r are non-negative integers, then $p > q$ and $a \in \{\frac{2-\frac{4}{n}}{s}, \frac{1-\frac{4}{n}}{s}\}$ for some positive integer s .

Conjecture 7. For any integer $n \geq 5, n \neq 28$ if $a \in \{\frac{1}{4}, \frac{1}{5}, \frac{2}{5}, \frac{3}{7}, \frac{1}{3}\}$ and $a \in \{\frac{2-\frac{4}{n}}{s}, \frac{1-\frac{4}{n}}{s}\}$ for some positive integer s , then either a or $1 - a$ is in $\{\frac{2}{n}, \frac{4}{n}, \frac{1}{3} + \frac{4}{3n}\}$.

Sketch of proof of conjecture 1. Suppose to the contrary there is a tiling of a regular n -gon with right triangles of angles $\alpha \leq \frac{\pi}{4}$, $\frac{\pi}{2} - \alpha$ and $\frac{\pi}{2}$, and $\alpha \notin \{\frac{\pi}{n}, \frac{2\pi}{n}, \frac{\pi}{6} + \frac{2\pi}{3n}\}$. Denote $a = \frac{2\alpha}{\pi}$.

If $\alpha = \frac{\pi}{4}$ then the angle of regular n -gon should be a rational multiple of $\frac{\pi}{4}$, hence only regular 4, 8 - gons can be tiled with right isosceles triangles. For $n = 8$, $\alpha = \frac{2\pi}{8} = \frac{\pi}{4}$.

Take a vertex of the n -gon. By p, q, r we denote the number of smaller acute angles α , bigger acute angles $\frac{\pi}{2} - \alpha$ and angles $\frac{\pi}{2}$, respectively at this vertex. Then $2 - \frac{4}{n} = pa + q(1 - a) + r$, for some $0 < a \leq \frac{1}{2}$. Then by conjecture 6, $p > q$ and $\alpha \in \{\frac{\pi-\frac{2\pi}{n}}{s}, \frac{\frac{\pi}{2}-\frac{2\pi}{n}}{s}\}$ for some positive integer s .

Since $\alpha \notin \{\frac{\pi}{n}, \frac{2\pi}{n}, \frac{\pi}{6} + \frac{2\pi}{3n}\}$, it follows that by conjecture 7, $\alpha \notin \{\frac{\pi}{10}, \frac{\pi}{5}, \frac{3\pi}{14}, \frac{\pi}{6}, \frac{\pi}{8}\}$.

For the triangles that have same vertices inside of regular n -gon not on the side of a triangle, by p we denote the number of the smaller acute angles α , by q the number of greater acute angles $\frac{\pi}{2} - \alpha$, by r the number of right angles. Then by conjecture 5 at any point inside of n -gon not on the side of a triangle $p \geq q$.

For the triangles that have same vertices on the side of n -gon or on the side of a triangle, we use conjecture 5, substituting $r - 2$ for r . Then at any point on the side of n -gon or on the side of a triangle $p \geq q$.

Hence the number of smaller acute angles is greater than the number of bigger acute angles. A contradiction. \square

Sketch of proof of conjecture 6. Assume to the contrary that $q - p \geq 0$.

We have

$$2 - \frac{4}{n} = pa + q(1 - a) + r = (q - p)(1 - a) + p + r.$$

If $q - p = 0$ then $2 - \frac{4}{n} = p + r$, but $2 > 2 - \frac{4}{n} > 1$ and $p + r$ is integer. Then $q - p \geq 1$.

If $r + p > 1$, then $1 - a = \frac{2 - \frac{4}{n} - p - r}{q - p} < 0$. A contradiction. Thus either $r + p = 1$ or $r + p = 0$.

If $r + p = 1$, then $1 - a = \frac{1 - \frac{4}{n}}{q - p}$. Hence $a = \frac{4}{n}$.

If $r + p = 0$, then $1 - a = \frac{2 - \frac{4}{n}}{q - p}$. Hence $q - p \in \{2, 3\}$. Then $a \in \{\frac{2}{n}, \frac{1}{3} + \frac{4}{3n}\}$. A contradiction. Thus $p > q$.

We have

$$2 - \frac{4}{n} = pa + q(1 - a) + r = (p - q)a + q + r.$$

Hence $0 \leq r + q < 2$.

Let $u = q + r$ and $s = p - q$. Then $a = \frac{2 - u - \frac{4}{n}}{s}$, where s is a non-negative integer.

Thus $a \in \{\frac{2 - \frac{4}{n}}{s}, \frac{1 - \frac{4}{n}}{s}\}$ for some positive integer s .

□

Sketch of proof of conjecture 7. We provide proof for general $a = \frac{z - \frac{4}{n}}{s}$ where z is either 1 or 2.

Suppose that $a = \frac{1}{4} = \frac{z - \frac{4}{n}}{s}$. Hence $s = 4z - \frac{16}{n}$. Since s and n are positive integers, it follows that $n \in \{1, 2, 4, 8, 16\}$. But $n \geq 5$ and $a = \frac{1}{4} = \frac{2}{8} = \frac{4}{16}$. Thus for $a = \frac{1}{4}$, $a \in \{\frac{2}{n}, \frac{4}{n}, \frac{1}{3} + \frac{4}{3n}\}$ for some positive integer n .

Suppose that $a = \frac{1}{5} = \frac{z - \frac{4}{n}}{s}$. Hence $s = 5z - \frac{20}{n}$. Since s and n are positive integers, it follows that $n \in \{1, 2, 4, 5, 10, 20\}$. But $n \geq 5$ and $a = \frac{1}{5} = 1 - \frac{4}{5} = \frac{2}{10} = \frac{4}{20}$. Thus for $a = \frac{1}{5}$, $a \in \{\frac{2}{n}, \frac{4}{n}, \frac{1}{3} + \frac{4}{3n}\}$ for some positive integer n .

If $a = \frac{2}{5} = \frac{z - \frac{4}{n}}{s}$. Hence $2s = 5z - \frac{20}{n}$. Since s and n are positive integers, it follows that $n \in \{1, 2, 4, 5, 10, 20\}$. But $n \geq 5$ and $a = \frac{2}{5} = \frac{2}{5} = \frac{4}{10} = \frac{1}{3} + \frac{4}{3 \cdot 20}$. Thus for $a = \frac{2}{5}$, $a \in \{\frac{2}{n}, \frac{4}{n}, \frac{1}{3} + \frac{4}{3n}\}$ for some positive integer n .

If $a = \frac{3}{7} = \frac{z - \frac{4}{n}}{s}$. Hence $3s = 7z - \frac{28}{n}$. Since s and n are positive integers, it follows that $n \in \{1, 2, 4, 7, 14, 28\}$. But $n \geq 5$, $n \neq 28$ and $a = \frac{3}{7} = \frac{1}{3} + \frac{4}{3 \cdot 14} = 1 - \frac{4}{7}$. Thus for, $a = \frac{3}{7}$, $a \in \{\frac{2}{n}, \frac{4}{n}, \frac{1}{3} + \frac{4}{3n}\}$ for some positive integer n .

If $a = \frac{1}{3} = \frac{z - \frac{4}{n}}{s}$. Hence $s = 3z - \frac{12}{n}$. Since s and n are positive integers, it follows that $n \in \{1, 2, 3, 4, 6, 12\}$. But $n \geq 5$ and $a = \frac{1}{3} = \frac{2}{6} = \frac{4}{12}$. Thus for $a = \frac{1}{3}$, $a \in \{\frac{2}{n}, \frac{4}{n}, \frac{1}{3} + \frac{4}{3n}\}$ for some positive integer n .

□

Remark 8. (*Relation to earlier results*)

(a) A tiling of a regular n -gon with right non-isosceles triangles with angle $\alpha \neq \frac{\pi}{4}$ is called regular if either

(1) at each point of the (convex hull of the) n -gon the number of angles α is equal to the number of angles $\frac{\pi}{2} - \alpha$, or

(2) at each point of the n -gon the number of angles α is equal to the number of angles $\frac{\pi}{2}$, or

(3) at each point of the n -gon the number of angles $\frac{\pi}{2} - \alpha$ is equal to the number of angles $\frac{\pi}{2}$.

Otherwise the tiling is called irregular.

(b) From [L12, Theorem 2.1] it follows that a regular n -gon, $n \geq 5, n \neq 6$ cannot be regularly tiled with congruent right triangles .

(c) Let us present a simple deduction of (b) from [L12, Theorem 2.1]. If a regular n -gon, $n \geq 5, n \neq 6$, is regularly tiled with triangles with angles $\alpha, \frac{\pi}{2} - \alpha, \frac{\pi}{2}$ then one of the properties (1), (2), (3) of (a) is true. Since $n \geq 5, n \neq 6$ cases (i, iv, vi, ix) listed in [L12, Theorem 2.1] cannot be true. Since the n -gon is regular and triangles are right, cases (ii, vii, viii) cannot be true.

In case (iii) we have either $\alpha = \frac{2\pi}{n}$ or $\frac{\pi}{2} - \alpha = \frac{2\pi}{n}$ and $\cos \alpha, \cos(\frac{\pi}{2} - \alpha)$ should be rational. Since $n \notin \{1, 2, 3, 4, 6\}$, it follows that $\cos(\frac{2\pi}{n})$ is not rational, according to [M, Theorem 4.1]. Hence case (iii) cannot be true.

In case (v) we have either $\alpha = \frac{2\pi}{n}$ or $\frac{\pi}{2} - \alpha = \frac{2\pi}{n}$ and $\frac{\sin \alpha}{\sin(\frac{\pi}{2} - \alpha)} = 1$. Thus $\alpha = \frac{\pi}{2} - \alpha = \frac{\pi}{4}$, but $n > 4$. Hence case (v) cannot be true.

(d) In [L20, Theorem 1] we have the assumption that, $n \geq 25, n \neq 30, 42$. In [L12, Theorem 2.1] we have the assumption that the tiling is regular, we do not know if for any $5 \leq n \leq 24, n \in 30, 42$ there are irregular tilings with right triangles with angle $\alpha \neq \frac{\pi}{n}$.

(e) For the cases $n < 5$ and $n = 6$ see [L12, Theorem 2.1] and [S, Theorem].

[L12] M.Laczkovich, Tilings of Convex Polygons with Congruent Triangles, Discrete Comput. Geom. 48, (2012) 330-372

[L20] M.Laczkovich, Irregular tilings of regular polygons with similar triangles, <https://arxiv.org/abs/2002.12013>

[M] A. Skopenkov. Mathematics Through Problems: from olympiades and math circles to profession. Part I. Algebra. AMS, Providence, https://www.mccme.ru/circles/oim/algebra_eng.pdf

[S] B. Szegedy, Tilings of the square with similar right triangles, Combinatorica 21 (1) (2001) 139-144

[V] L.Vigdorichik, Tiling of regular polygon with similar right triangles II, <https://www.mccme.ru/circles/oim/mmks/works2020/vigdorichik1.pdf>